

Revised in February, 1997

The outer derivation of a complex Poisson manifold.

by J-L. Brylinski* and G. Zuckerman

In this note we introduce the canonical outer Poisson vector field on a Poisson manifold (both in the smooth and complex-analytic cases). This means that we construct a canonical class of Poisson vector fields modulo the locally hamiltonian vector fields. This class is an obstruction to the existence of a volume form invariant under all hamiltonian flows. It is defined even both in the case of a smooth Poisson manifold and in the case of a holomorphic Poisson manifold, even if there is no global volume form. We show (in the complex-analytic case) that this outer vector field gives a class of holomorphic functions with singularities along the complement of the regular set, which is canonical modulo the addition of Casimir functions. We study this class in terms of various sheaf cohomology groups of interest. The same outer vector field has been introduced independently by Alan Weinstein [W]. He has a description of the outer vector field which relates it to the KMS theory and thus to the Connes-Takesaki theory of modular automorphisms of a von Neumann algebra.

We thank Alan Weinstein for communicating to us an early version of his work and for useful correspondence. The first author thanks the participants at the Singularity Seminar at Marseille-Château Gombert where he lectured on this work. We thank Philip Foth for useful comments on a previous version of this note.

1. Hamiltonian and outer vector fields of a Poisson manifold

We will work either with a C^∞ or a complex Poisson manifold M . In the first case there is a Poisson tensor $\pi \in \wedge^2 TM$, where TM is the tangent bundle. In the second case the Poisson tensor π is a holomorphic section of $\wedge^2 \Theta_M$, where Θ_M is the holomorphic tangent bundle.

There are three interesting classes of vector fields, which we enumerate starting with the largest class:

(1) the *Poisson vector fields*: a vector field ξ is Poisson if it preserves the Poisson structure, that is

$$\mathcal{L}_\xi \pi = \{\xi, \pi\} = 0,$$

where $\{ , \}$ is the Schouten bracket.

(2) the *locally hamiltonian vector fields*: locally ξ is of the form $X_H = i(dH)\pi$, where i denotes interior product, and H is a smooth (resp. holomorphic) function defined locally.

(3) the hamiltonian vector fields: $\xi = X_H$ for some global H .

* This research was supported in part by NSF grant DMS-9504522.

It is interesting to devise cohomological criteria separating these types. For instance, let ξ be locally hamiltonian; then there is an open covering (U_i) such that $\xi = X_{H_i}$ over U_i . Then $f_{ij} = H_j - H_i$ satisfies $X_{f_{ij}} = 0$, so it is a Casimir function over U_{ij} . We have a Čech 1-cocycle f_{ij} with values in the sheaf $\mathcal{C}_M$ of smooth (resp. holomorphic) Casimir functions.

Proposition 1. (1) For M a smooth Poisson manifold, we have an exact sequence

$$0 \rightarrow \text{Ham}(M) \rightarrow \text{Loc Ham}(M) \rightarrow H^1(M, \mathcal{C}_M) \rightarrow 0.$$

(2) For M a complex Poisson manifold, we have an exact sequence

$$0 \rightarrow \text{Ham}(M) \rightarrow \text{Loc Ham}(M) \rightarrow H^1(M, \mathcal{C}_M) \rightarrow H^1(M, \mathcal{O}_M),$$

where $\mathcal{O}_M$ is the sheaf of germs of holomorphic functions.

Now we wish to examine the difference between the first two types of vector fields. We need to sheafify the question. Let $\underline{\text{Ham}}$ be the sheaf of locally hamiltonian vector fields, and let $\underline{\text{Der}}$ be the sheaf of Poisson vector fields. Then $\underline{\text{Ham}}$ is a subsheaf of $\underline{\text{Der}}$. If we have a vector field ξ which is Poisson, it gives a section of the quotient sheaf $\underline{\text{Der}}/\underline{\text{Ham}}$. This quotient sheaf is hard to describe in general. In the regular case we have the following description, where $\underline{TX}$ denotes the sheaf of smooth tangent vectors, $\underline{\Theta}_X$ the sheaf of holomorphic tangent vectors.

Proposition 2. Let M be a regular Poisson manifold such that there exists a smooth fibration $p : M \rightarrow X$ whose fibers are the symplectic leaves. Then the quotient sheaf $\underline{\text{Der}}/\underline{\text{Ham}}$ identifies with the pull-back sheaf $p^{-1}\underline{TX}$ (in the smooth case), resp. $p^{-1}\underline{\Theta}_X$ (in the holomorphic case).

Now in the smooth case the sheaf cohomology $H^i(M, p^{-1}\underline{TX})$ is equal to $\text{Vect}(X) \otimes H^i(F, \mathbb{R})$, where F is the fiber of p . In the holomorphic case, if M is Stein, the group $H^i(M, p^{-1}\underline{\Theta}_X)$ is equal to $\Theta(X) \otimes H^i(F, \mathbb{C})$. In general there is a spectral sequence with E_2 term $E_2^{pq} = H^p(X, \underline{\Theta}_X \otimes H^q(F, \mathbb{C}))$ converging to $H^i(M, p^{-1}\underline{\Theta}_X)$. So we obtain

Proposition 3. Let M be as in Proposition 2. Let ξ be a Poisson vector field on M . Then ξ is locally hamiltonian iff the corresponding vector field on X is trivial.

Now we come to the notion of an outer Poisson vector field. This is defined to be a section of the quotient sheaf $\underline{\text{Der}}/\underline{\text{Ham}}$.

Lemma 1. Let M be a smooth Poisson manifold or a complex Poisson manifold which is Stein. An outer Poisson vector field comes from a global Poisson vector field iff the corresponding class in $H^1(M, \underline{\text{Ham}}) \xrightarrow{\sim} H^2(M, \mathcal{C}_M)$ is trivial.

The isomorphism comes from the exact sequences of sheaves

$$0 \rightarrow \mathcal{C}_M \rightarrow \underline{\mathcal{C}}_M^\infty \rightarrow \underline{\text{Ham}}_M \rightarrow 0,$$

in the smooth case

$$0 \rightarrow \mathcal{C}_M \rightarrow \mathcal{O}_M \rightarrow \underline{Ham}_M \rightarrow 0,$$

in the holomorphic case. In the holomorphic case we have $H^1(M, \mathcal{O}_M) = H^2(M, \mathcal{O}_M) = 0$ because M is Stein. ■

2. Koszul operators and the canonical outer vector field.

The results here were also obtained independently by A. Weinstein [W]. Following Koszul [K], we say that an operator $D : \wedge^i TM \rightarrow \wedge^{i-1} TM$ generates the Schouten bracket iff we have

$$\{u, v\} = (-1)^p [D(u \wedge v) - (Du) \wedge v - (-1)^p u \wedge (Dv)], \quad (1)$$

where $u \in \wedge^p TM$, $v \in \wedge^q TM$. There is a similar notion in the holomorphic case. We call these operators Koszul operators.

Theorem 1. (Koszul) (1) *The set of Koszul operators D is in canonical bijection with the set of connections ∇ on the canonical bundle $\wedge^n T^*M$ (resp. $\omega_M = \underline{\Omega}_M^n$).*

(2) *The square D^2 is interior product with the curvature K of ∇ .*

(3) *Let ∇ be a connection on the canonical bundle, and let α be a 1-form. Then the Koszul operator corresponding to the connection $\nabla + \alpha$ is $D + i(\alpha)$.*

In particular, a Koszul operator is of square 0 iff the corresponding connection is flat.

Recall that a holomorphic vector bundle $E \rightarrow M$ has Chern classes $c_p(E) \in H^p(M, \underline{\Omega}^p)$ in Hodge cohomology, and slightly more refined classes in $H^p(M, \underline{\Omega}_{M,cl}^p)$, where $\underline{\Omega}_{M,cl}^p$ is the sheaf of closed holomorphic p -forms.

Corollary . *Let M be a complex manifold.*

(1) *There exists a global holomorphic Koszul operator iff the first Chern class $c_1(\Theta_M) \in H^1(M, \underline{\Omega}_M^1)$ of Θ_M in Hodge cohomology is 0.*

(2) *There exists a global holomorphic Koszul operator of square 0 iff the class $c_1(\Theta_M) \in H^1(M, \underline{\Omega}_{M,cl}^1)$ is 0.*

We recall how a connection ∇ defines a Koszul operator. We discuss only the holomorphic case. We have the canonical isomorphism

$$* : \wedge^i \Theta_M \xrightarrow{\sim} \underline{\Omega}_M^{n-i} \otimes \omega_M^{\otimes -1},$$

where $\underline{\Omega}_M^i$ is the sheaf of holomorphic i -forms. The connection ∇ on ω_M induces one on the dual line bundle $\omega_M^{\otimes -1}$, which in turn extends to operators $\underline{\Omega}_M^j \otimes \omega_M^{\otimes -1} \rightarrow \underline{\Omega}_M^{j+1} \otimes \omega_M^{\otimes -1}$. Using $*$ this induces an operator $D : \wedge^i \Theta_M \rightarrow \wedge^{i-1} \Theta_M$. It is easy to see that D is a Koszul operator.

Note that any volume form ν induces a flat connection ∇ on ω_M , hence a Koszul operator of square 0 which we will denote by D_∇ . More generally any multi-valued volume form ν , such that monodromy around any loop multiplies ν by some constant, also defines a flat connection.

Now assume that M is a Poisson manifold with Poisson tensor π . Then for any (local) Koszul operator D of square 0, the vector field $D\pi$ is Poisson because (1) gives

$$0 = D\{\pi, \pi\} = -(D\pi) \wedge \pi - \pi \wedge D\pi = -2(D\pi) \wedge \pi$$

and then

$$2\{D\pi, \pi\} = 2D^2(\pi) \wedge \pi - D^2(\wedge^2 \pi) = 0$$

(cf. [K]).

Now if we change the Koszul operator of square 0 to $D + i(df)$, then we change $D\pi$ to $D\pi + X_f$. Hence we have

Theorem 2. *There exists a canonical section ξ of $\underline{Der}/\underline{Ham}$, which is locally equal to $D\pi$ for any Koszul operator D of square 0.*

This is called the *canonical outer vector field*.

The canonical outer vector field can be computed more concretely from any local volume form ν . We have

Proposition 4. *Given a volume form ν over $U \subseteq M$, a representative of ξ over U is given by the vector field v such that $(v \cdot f)\nu = \mathcal{L}_{X_f}\nu$.*

Let us now look at the regular case. Then we have

Proposition 5. *For a regular Poisson manifold equipped with a fibration $p : M \rightarrow Z$ whose fibers are the symplectic leaves, and for any volume form ν , the Poisson vector field $D\pi$ is equal to*

$$\pm X_{\log f},$$

where the function f is defined as follows. Let α be some volume form on Z ; then the Liouville fiberwise volume form β defines a volume form $\alpha \wedge \beta$ on M . Then

$$f = \frac{\alpha \wedge \beta}{\nu}.$$

Corollary . (see also [W]) *The section of $\underline{Der}/\underline{Ham}$ is supported in the singular set of the Poisson manifold M .*

As an example, let $\mathfrak{g}$ be a complex Lie algebra, and let $\lambda : \mathfrak{g} \rightarrow \mathbb{C}$ be the modular character $\lambda(\xi) = \text{Tr}(ad(\xi))$. Then the constant vector field λ on $\mathfrak{g}^*$ represents the outer vector field ξ . We have

Proposition 6. *Let $\mathfrak{g}$ be a (real or complex) Lie algebra which is not unimodular. Then the outer Poisson vector field ξ does not vanish in any neighborhood of 0.*

This is proved easily by looking at the Taylor series at the origin of any hamiltonian vector field.

3. The canonical singularity class.

For a Poisson manifold M , we have the canonical Poisson outer vector field ξ . We have proved that its restriction to the regular open set U is trivial. We wish to think of ξ as corresponding to a class (modulo Casimir functions) of singularities along $S = M \setminus U$ of multivalued meromorphic functions. This will be mostly interesting in the complex-analytic case. We use cohomology $H_S^i(M, F)$ with supports in S of a sheaf F , which fits into the exact sequence

$$0 \rightarrow H_S^0(M, F) \rightarrow H^0(M, F) \rightarrow H^0(U, F) \xrightarrow{\partial} H_S^1(M, F) \rightarrow H^1(M, F) \rightarrow \dots$$

Proposition 7. (1) *Assume that M is equipped with a holomorphic volume form ν , and let D be the corresponding Koszul operator. Then the Poisson vector field $v = D\pi$ is locally hamiltonian over U , hence it gives rise to a class $\bar{f}_\nu$ in $H^0(U, \mathcal{O}_M/\mathcal{C}_M)$.*

(2) *The outer vector field ξ gives a canonical class in $H_S^1(M, \mathcal{O}_M/\mathcal{C}_M)$. If there is a global volume form, this class is the image of the class $\bar{f}_\nu$ in (1) by the boundary operator ∂ .*

Here $H_S^1(M, \mathcal{O}_M/\mathcal{C}_M)$ denote sheaf cohomology with supports in S .

In (1) we could more generally assume that M has a multi-valued volume form ν such that $\frac{T\nu}{\nu}$ is constant for any monodromy transformation T .

This class can be understood as follows. First assume there is a global volume form. The global Poisson vector field $v = D\pi$ which over U is locally hamiltonian, hence given by a section of $\mathcal{O}_M/\mathcal{C}_M$ over U . We can change the volume form, multiplying it by a non-vanishing function f over M . Then $\log f$ is globally defined over U modulo constants, hence it is globally defined as a section of the quotient sheaf $\mathcal{O}_M/\mathcal{C}_M$ over U . Therefore the image in $H_S^1(M, \mathcal{O}_M/\mathcal{C}_M)$ is well-defined.

Now relax the assumption about a global volume form. The cohomology with supports $H_S^*(M, \mathcal{O}_M/\mathcal{C}_M)$ can be computed as a direct limit over open coverings (U_i) of M of the quotient complex $C^\bullet((U_i \setminus (S \cap U_i), F)/C^\bullet((U_i), F))$ where $F = \mathcal{O}_M/\mathcal{C}_M$. This is a complex whose $(p+1)$ -st term is

$$\oplus_{i_0, \dots, i_p} \Gamma(U_{i_0 \dots i_p} \setminus (S \cap U_{i_0 \dots i_p}), F) / \Gamma(U_{i_0 \dots i_p}, F).$$

This description is correct because there is no section of F on an open set which vanishes on a dense open subset.

Now cover S by open sets $S \cap U_i$, U_i open in M , such that there is a volume form ν_i over U_i . Then we have a corresponding Poisson vector field v_i over U_i ; the

restriction of v_i to $U_i \setminus (S \cap U_i)$ is locally hamiltonian, hence defines a section g_i of $\underline{Ham} \xrightarrow{\sim} \mathcal{O}_M/\mathcal{C}_M$ over this open set. Now we have: $v_j - v_i = X_{\log \frac{\nu_j}{\nu_i}}$ over U_{ij} hence if we put $f_{ij} = \log \frac{\nu_j}{\nu_i}$, we get a section of $\mathcal{O}_M/\mathcal{C}_M$ over U_{ij} such that $v_j - v_i = X_{f_{ij}}$, hence $g_j - g_i = f_{ij}$. It follows that the Čech coboundary of (g_i) is zero in the quotient complex, hence we have indeed a cohomology class in degree 1.

There is another interesting cohomological invariant attached to ξ . This is the class in $H_S^2(M, \mathcal{C}_M)$ obtained from the class in $H_S^1(M, \mathcal{O}_M/\mathcal{C}_M)$ by applying the boundary map for the exact sequence of sheaves $0 \rightarrow \mathcal{C}_M \rightarrow \mathcal{O}_M \rightarrow \mathcal{O}_M/\mathcal{C}_M \rightarrow 0$.

If we have a volume form on M , then there is a corresponding class in $H^1(U, \mathcal{C}_M)$. We can summarize the situation into the following square

$$\begin{array}{ccc} H^0(U, \mathcal{O}_M/\mathcal{C}_M) & \xrightarrow{\partial} & H^1(U, \mathcal{C}_M) \\ \downarrow \partial & & \downarrow \partial \\ H_S^1(M, \mathcal{O}_M/\mathcal{C}_M) & \xrightarrow{\partial} & H_S^2(M, \mathcal{C}_M). \end{array}$$

The classes in the groups of the first row depend on a volume form; the classes in the groups in the second row are entirely intrinsic.

We would like to suggest that the groups $H^1(U, \mathcal{C}_M)$ and $H_S^2(M, \mathcal{C}_M)$ have a geometric interpretation in terms of multivalued holomorphic functions on U whose variation (under monodromy) is a Casimir function. We first study the case where the Poisson tensor π is non-degenerate on a dense open set U . Then the vanishing locus of the section $\wedge^n \pi$ of $\wedge^n \Theta_X$ is a divisor $D = \sum_i n_i D_i$. The singular locus S is the support of D , i.e., the union of the D_i . Since all the Casimir functions on any open set are constant, the sheaf $\mathcal{C}_M$ is the constant sheaf $\underline{\mathbb{C}}_M$. We have $H_S^2(M, \mathbb{C}) = \sum_i \mathbb{C}[D_i]$, where D_i is the class of the divisor in cohomology with support. Then we can state

Proposition 8. *The class in $H_S^2(M, \mathbb{C})$ corresponding to the canonical outer vector field is the class $\sum_i n_i [D_i]$ of the divisor of $\wedge^n \pi$.*

We illustrate this in a simple example. Let $\mathfrak{g}$ be the two-dimensional complex Lie algebra with basis x, y , such that $[x, y] = x$. Then $\mathfrak{g}^* = \mathbb{C}^2$ is a holomorphic Poisson manifold, with Poisson tensor $\pi = x \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y}$. The regular open set U is the open set $U = \{(x, y); x \neq 0\}$. The singular set S is the line $x = 0$. If we use the volume form $dx \wedge dy$, then we find that ξ is represented by the Poisson vector field $v = \frac{\partial}{\partial y}$. Over a contractible open set in U , this is the hamiltonian vector field associated to a branch $\log x$ of the logarithm function. The obstruction to finding a global function f such that $v = X_f$ is given by the monodromy of $\log x$, which is encoded in the generator of $H^1(U, \mathbb{C}) = \mathbb{C}$. The class in $H_S^2(M, \mathbb{C}) = \mathbb{C}$ is the canonical orientation class.

There is a similar computation for quadratic Poisson algebras, and for the Poisson-Lie group corresponding to $SL(2)$.

In general the group $H^1(U, \mathcal{C}_M)$ is hard to describe. However for any symplectic leaf $F \subset U$ we have a restriction map $H^1(U, \mathcal{C}_M) \rightarrow H^1(F, \mathbb{C})$; therefore for any loop $\gamma \in F$ we have a corresponding character $\chi_\gamma : H^1(U, \mathcal{C}_M) \rightarrow \mathbb{C}$. This has the following description:

Proposition 9. *Let U be an open subset of M over which there is a fibration $p : M \rightarrow Z$ whose fibers are symplectic leaves. Assume there is a multi-valued global volume form ν on U , and let $c \in H^1(U, \mathcal{C}_M)$ be the corresponding cohomology class. Let $F \subset U$ be a symplectic leaf, and let $x = p(F)$. Let α be an element of $\wedge^{\max} \Theta_{X,x}^*$, and let g be the multi-valued function on F given by*

$$g = \frac{\alpha \wedge \beta}{\nu},$$

where β is the fiberwise volume form as in Proposition 5. Then for any loop $\gamma \in F$ the number $\chi(c)$ is equal to the variation of the multi-valued function $\log g$ along the loop γ .

Concerning the cohomology groups $H^i(M, \mathcal{C}_M)$ associated to a complex Poisson manifold M , they are modules over the algebra $H^0(M, \mathcal{C}_M)$ of global Casimir functions. It is tempting to conjecture that for M a Stein manifold, these are finitely-presented modules.

References:

[K] J-L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie*, Astérisque vol. **Hors-Série**: Elie Cartan et les Mathématiques d'Aujourd'hui, Soc. Math. Fr. (1985), 257-271

[W] A. Weinstein, *The modular automorphism group of a Poisson manifold*, preprint (1995)